



# Exploring Psychosocial Experiences through a Phenomenological Qualitative Study of Using mHealth Applications Among Pregnant Women in Rural Communities

Sarce Makaba <sup>1\*</sup>, Helmin Rumbiak <sup>2</sup>

<sup>1</sup>Universitas Cenderawasih, Indonesia

<sup>2</sup>Universitas Cenderawasih, Indonesia

<sup>1</sup>[sarcemakaba@gmail.com](mailto:sarcemakaba@gmail.com), <sup>2</sup> [rumbiakhelmin@gmail.com](mailto:rumbiakhelmin@gmail.com)

## Article Info

### Article history:

Received 27-07-2025

Revised 19-09-2025

Accepted 24-09-2025

### Keyword:

Psychosocial Experiences, mHealth Applications, Pregnant Women, Rural Communities, Digital Health, Qualitative Study

## ABSTRACT

Digital health technologies have become an integral part of maternal healthcare, particularly through the growing use of mobile health (mHealth) applications to support pregnancy monitoring. While mHealth offers access and information benefits, little is known about how pregnant women in rural communities emotionally and socially experience these tools in their daily lives. Existing studies have focused on usage patterns and clinical outcomes, yet they have not explored the deeper psychosocial meanings that shape engagement with mHealth—raising the question: how do pregnant women in underserved areas experience and interpret the use of mHealth applications during pregnancy? This study uses a descriptive phenomenological approach to explore the lived experiences of pregnant women using mHealth tools in rural settings. Participants consisted of twelve pregnant women aged 19–36 years, most of whom were in their second and third trimesters, with varied parity status. All participants resided in geographically remote rural villages characterized by limited internet infrastructure, restricted access to specialized healthcare, and socioeconomic challenges that constrained consistent digital use. Data were collected through in-depth semi-structured interviews and analyzed using Colaizzi's method to extract essential themes. The analysis revealed four core experiential themes: emotional ambivalence, digital isolation, cognitive fatigue, and evolving trust. These findings demonstrate that mHealth engagement is influenced not only by functional aspects but also by emotional readiness, social support, and the usability of digital tools in contextually limited environments. This study contributes to a more holistic understanding of digital health experiences and calls for future research that centers users' voices to develop more inclusive and sustainable mHealth solutions.



©2025 Authors. Published by PT Mukhlisina Revolution Center.. This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

(<https://creativecommons.org/licenses/by/4.0/>)

## INTRODUCTION

The integration of digital health technologies into maternal care has emerged as a global response to the growing demand for accessible, personalized, and cost-effective health services (Antelo et al., 2022). Among these innovations, mobile health (mHealth) applications have become increasingly prevalent in supporting pregnancy monitoring, offering tools such as symptom tracking, health education, appointment reminders, and remote consultations (Rabbitts et al., 2021). These platforms are particularly promoted in underserved and rural areas where access to conventional healthcare services remains limited. In many low- and middle-income countries, including those with vast rural populations, mHealth is positioned as a potential bridge to close healthcare gaps and empower expectant mothers with timely information and guidance.

Despite the technological promise of mHealth systems, the lived realities of their users—especially pregnant women in marginalized communities—are often underexplored (Brogly et al.,

2021). For many women, the experience of pregnancy is deeply influenced by cultural expectations, familial responsibilities, and emotional fluctuations. Introducing digital tools into this intimate life event can either enhance or disrupt the perceived sense of security, agency, and connection to care (Emert et al., 2023). This highlights the importance of moving beyond functional evaluations to understand how women emotionally and socially engage with mHealth in daily life.

Understanding how women experience and make meaning of these technologies during pregnancy is crucial. Their perspectives offer more than functional feedback; they reveal the emotional, psychological, and cultural dimensions embedded in the interaction between user and technology (van der Storm, Bemelman, et al., 2023). Yet, Current evaluations of mHealth interventions tend to emphasize measurable outcomes—such as adherence, clinical indicators, or usability—while overlooking the subjective meanings that influence trust and sustained engagement (Potzel et al., 2021). This research addresses that gap by exploring the psychosocial dimensions of pregnant women’s experiences with mHealth in rural contexts.

Adopting a phenomenological approach provides a pathway to reveal these nuanced experiences, capturing the emotional, psychological, and cultural dimensions often invisible in quantitative studies (Wan et al., 2021). Rather than assuming uniform acceptance, this perspective foregrounds the complexity of how individuals perceive and live through digital maternal care, especially in socially and digitally constrained environments.

In recent years, research on individuals’ lived experiences in health-related contexts has gained increasing prominence, particularly as scholars and practitioners seek to understand how users engage with digital health interventions beyond technical functionality (van der Storm et al., 2024). By centering the voices of pregnant women, this study contributes novel insights into how mHealth technologies are internalized during pregnancy, thereby extending the discourse from technical utility toward a more holistic understanding of digital maternal health (Pickard et al., 2025). These experiences are deeply personal, shaped by individual beliefs, cultural frameworks, emotional states, and varying levels of digital literacy.

However, capturing the depth of such experiences poses significant methodological challenges (Sirbu et al., 2025). Most existing studies on mHealth adoption and maternal health outcomes rely heavily on quantitative metrics—such as app usage frequency, patient adherence rates, or outcome-based indicators—that, while valuable, fall short in representing the emotional, psychological, and cultural nuances embedded in users’ experiences (Ben-Zeev et al., 2024). Surveys and structured assessments often overlook how users interpret and emotionally respond to technology, especially when operating in environments with social and infrastructural constraints.

Consequently, many prior approaches have lacked the sensitivity and depth needed to uncover the essential meanings that participants assign to their interaction with mHealth tools (Badawy et al., 2021). Without qualitative insight, particularly one grounded in phenomenology, these studies risk offering partial or superficial interpretations of the user experience (Psihogios et al., 2021). This limitation not only restricts theoretical understanding but also hampers the development of responsive, culturally informed digital health solutions.

A descriptive phenomenological approach, therefore, becomes essential in filling this gap (Wong et al., 2020). By centering on the subjective accounts of pregnant women and capturing the structures of their lived experience, such research provides access to meanings that are otherwise invisible in data-driven evaluations (King et al., 2023). It allows for a more holistic understanding of how technology is received, internalized, and lived in the everyday reality of pregnancy in rural communities.

Current efforts to enhance maternal health in rural areas have primarily emphasized practical interventions, such as implementing mHealth platforms to improve access to antenatal care, streamline communication with health providers, and deliver timely health information (Birrell et al., 2022). These solutions have shown promise in addressing logistical and service-related challenges (Sauer et al., 2024). However, most evaluations of these interventions have employed quantitative

frameworks or outcome-driven models that emphasize efficiency and usage statistics over human-centered understanding.

While practical in design, these approaches are limited in their capacity to capture the subjective dimensions of technology use—particularly the emotional, psychological, and sociocultural complexities experienced by pregnant women navigating unfamiliar digital environments (Iglesias et al., 2023). Quantitative measures often fail to reveal the fears, uncertainties, and contextual realities that influence how mHealth is perceived, internalized, or even rejected by users in rural communities (Tatar et al., 2023). Consequently, the current knowledge landscape remains fragmented, with insufficient insight into the lived experiences that shape maternal engagement with digital health.

To develop more culturally responsive and emotionally attuned digital health strategies, there is a clear need for alternative approaches that center on human experience (Morton et al., 2022). Phenomenology, with its focus on describing the essence of lived phenomena as they are experienced by individuals, offers a powerful framework for exploring how pregnant women interpret and emotionally navigate the use of mHealth technologies (Fan et al., 2020). By privileging participants' voices and contextual realities, phenomenological inquiry allows for a more holistic and empathetic understanding of how digital tools intersect with maternal identity, vulnerability, and agency—dimensions that remain inadequately understood in current literature.

Previous studies have examined the adoption of mHealth in maternal care, mostly focusing on system usability, clinical outcomes, and access improvement. For example, Lattie et al. (2020) highlighted the functional benefits of mobile apps in low-income settings but did not explore users' emotional experiences. Giunti et al. (2025) identified social barriers to mHealth but did not offer in-depth insights into psychological impacts. These gaps show that the emotional and cultural experiences of pregnant women using mHealth remain under-researched. A deeper exploration is needed to understand how these technologies are lived and interpreted in everyday maternal life.

This study uses descriptive phenomenology to uncover the psychosocial experiences of pregnant women using mHealth applications in rural areas (Rangraz Jeddi et al., 2023). The method was chosen for its focus on lived meaning and its ability to describe how individuals perceive a phenomenon in their own terms (Cyriac et al., 2021). Colaizzi's approach was applied to organize and extract essential meanings from in-depth interview data. This approach answers the need identified in the previous section by providing rich, contextualized understanding of digital health engagement. It offers insight into the fears, hopes, and adaptations experienced by users in their unique environments.

The article is structured as follows: the introduction presents the problem and background. The next section explains the phenomenological framework and research context. This is followed by a description of the data collection and analysis process using Colaizzi's method. The results section presents major themes, supported by direct participant quotations. The article concludes with a discussion of the findings, their implications, and recommendations for digital maternal health design and policy.

## **RESEARCH METHODS**

### **Study Design**

A descriptive phenomenological design was employed to explore the psychosocial experiences of pregnant women using mobile health (mHealth) applications in rural settings (Norden et al., 2022). This approach was selected to allow for an in-depth examination of lived experiences, enabling a deeper understanding of the meanings individuals attribute to their interaction with digital health technologies during pregnancy. Rooted in Husserlian philosophy, descriptive phenomenology aims to uncover the essence of human experiences by bracketing preconceived notions and focusing solely on participants' descriptions (Mills, 2019). This design aligned with the study's objective to capture the nuanced emotional and cognitive dimensions of using mHealth platforms, particularly in underserved areas where technological access may be limited and culturally mediated.

## **Participants**

Participants consisted of twelve pregnant women residing in rural districts who had experience using mHealth applications for pregnancy monitoring. Selection followed a purposive sampling approach, with inclusion criteria encompassing women aged 18 years and older, currently in their second or third trimester, and having used any form of mobile health application consistently for at least four weeks. Exclusion criteria included individuals with diagnosed cognitive impairments or those who had discontinued use of the application due to unrelated medical issues. The participants' ages ranged from 22 to 39 years, with varied educational backgrounds and socioeconomic statuses. All were currently enrolled in local maternal health programs that incorporated digital monitoring tools. Data collection continued until no new themes or significant variations emerged during analysis, indicating that data saturation had been reached by the twelfth participant. Saturation was assessed iteratively, with the research team reviewing each successive transcript to determine whether additional interviews contributed novel insights or merely reinforced existing themes.

## **Data Collection**

Data were collected through in-depth, semi-structured interviews conducted face-to-face in a private setting, either in participants' homes or local community health posts. A flexible interview guide was utilized to encourage open narratives while ensuring relevance to the research focus. Interviews lasted between 45 and 75 minutes and were conducted in the participants' native language, then transcribed verbatim and translated into English for analysis. Comfortable and confidential environments were established to promote honest and reflective responses. All interviews were audio-recorded with participants' consent. The interview protocol was adapted from established phenomenological interview frameworks and piloted with two respondents prior to full data collection.

## **Data Analysis**

Data analysis was conducted using Colaizzi's descriptive phenomenological method. Transcripts were first read repeatedly to gain a holistic understanding (Johnson, 2014). Meaningful statements were then extracted and coded, followed by the formulation of meanings. These meanings were organized into clusters of themes that captured the essence of the experience. Thematic clusters were continually validated against the original transcripts to ensure fidelity to participants' expressions. NVivo software was used to facilitate the organization and retrieval of coded data without influencing interpretive processes. Essential structures of experience were synthesized into a comprehensive description that conveyed the psychosocial realities faced by the participants in their interaction with mHealth technologies.

## **Ethical Considerations**

Ethical approval for this study was obtained from the appropriate institutional review board prior to data collection. Informed written consent was obtained from all participants, with full disclosure of the study's aims, procedures, risks, and benefits. Confidentiality and anonymity were ensured by assigning pseudonyms and storing data in password-protected digital folders. The study adhered to the ethical principles outlined in the Declaration of Helsinki and complied with national guidelines for research involving human subjects.

## **RESULTS**

This section presents the lived experiences of twelve pregnant women in rural areas who used mobile health (mHealth) applications for pregnancy monitoring. Through Colaizzi's descriptive phenomenological method, four essential themes emerged from the participants' narratives. These themes reveal deep psychosocial dimensions of mHealth engagement, grounded in contextual, emotional, and technological realities.

### **Ambivalence Between Hope and Anxiety**

Participants expressed a duality of emotions—hopeful expectations from using digital tools for their pregnancy, contrasted with fear and insecurity in operating unfamiliar technology. While they acknowledged the potential benefits of monitoring fetal health and receiving timely information, this was often overshadowed by worry about misusing the application.

“I was happy when the midwife installed the app, but every time I had to input something, I felt scared—what if I press the wrong thing and harm my baby?” (P3)

“It’s supposed to help, but it made me more anxious. Sometimes I didn’t understand what the app was asking me.” (P7)

“I wanted to believe it was safe, but I kept asking myself, can a phone really tell me if my baby is okay?” (P11)

This ambivalence underscores a psychological tension that reveals a lack of digital confidence and a heightened sense of responsibility, particularly as it relates to the well-being of the unborn child.

### Balancing Hope and Fear in Digital Pregnancy Tools



### Digital Isolation and the Search for Emotional Support

Despite having access to digital maternal care, many participants described feeling emotionally isolated. The absence of human interaction in their use of mHealth heightened their sense of vulnerability and detachment, especially when they encountered difficulties in understanding instructions or interpreting app feedback.

“I missed having someone to talk to. The app is useful, but it doesn’t comfort you when you’re worried.” (P2)

“I still had to call the village midwife sometimes. The app can’t answer me like she can.” (P9)

“When I was worried about my baby’s movement, I opened the app, but it just showed numbers. I needed someone to explain, not just numbers.” (P8)

The digital divide here is not only infrastructural but emotional. The data suggests that participants often longed for empathetic reassurance—something technology alone could not provide in its current form.

### Technological Fatigue and Cognitive Overload

Participants frequently conveyed mental exhaustion stemming from the need to repeatedly engage with the application. Instructions were often perceived as complex, interfaces unfriendly, and notifications overwhelming. This led to avoidance behaviors and occasional disengagement.

“I felt tired just thinking about using it. Too many notifications. I didn’t even know which ones mattered.” (P6)

“Sometimes I didn’t open it for days. I knew I should, but it felt like a burden.” (P1)

“The words were too difficult. I had to ask my husband to read them, and that made me feel ashamed.” (P12)

This fatigue reveals a form of passive resistance to mHealth engagement, particularly when applications were not designed with local literacy, language preferences, and user capacity in mind.

### **Trust-Building Through Familiarization and Peer Influence**

While initial usage was marked by uncertainty, several participants developed greater confidence over time—especially those who received continuous guidance from peers or healthcare workers. Peer recommendation and community validation emerged as crucial in building trust in the app.

“At first, I didn’t trust it. But after my cousin explained and showed me how she used it, I started to feel okay.” (P10)

“The village health worker visited and showed me again how to check the baby’s growth. That helped me feel more sure.” (P4)

“When other mothers in the group said they also used it, I felt more confident. If they believed it worked, maybe I should too.” (P5)

This finding highlights the role of social learning and interpersonal interaction in overcoming technological resistance. Familiarity, when coupled with human facilitation, emerged as a critical factor in building sustained trust.

### **Essential Conclusion**

The psychosocial experience of using mHealth applications during pregnancy in rural contexts is shaped by a dynamic interplay of emotional vulnerability, digital alienation, cognitive burden, and interpersonal reassurance. These themes reveal that successful implementation of maternal digital health technologies must go beyond technical deployment, addressing the deeper human needs for understanding, emotional safety, and guided adaptation.

## **DISCUSSION**

The findings of this study reveal that the psychosocial experience of using mHealth during pregnancy is marked by emotional ambivalence, digital isolation, cognitive burden, and evolving trust (Atujuna et al., 2021). These essential themes respond directly to the central research question, which sought to understand how pregnant women in rural areas experience and assign meaning to their use of mobile health applications.

This study contributes a deeper understanding of how digital maternal health tools are lived and interpreted by users in socially and technologically constrained settings (Browne et al., 2020). Rather than assuming uniform usability, the findings uncover tensions between the promise of digital care and the psychological realities of using it. In doing so, this research adds nuance to the Technology Acceptance Model (TAM) by showing that acceptance is not solely a matter of perceived usefulness or ease of use, but also of emotional security, social reassurance, and cultural alignment. The themes illustrate that pregnant women do not merely adopt or reject mHealth based on technical functionality; instead, their engagement is shaped by emotions such as fear, reassurance, and fatigue, as well as the presence or absence of social support (Ward et al., 2022). This suggests that psychosocial readiness should be treated as a critical determinant in digital health adoption frameworks.

The findings resonate with prior research by Granholm et al. (2020), who acknowledged the value of mHealth for information access but did not examine users’ internal struggles. Similarly, Morcillo-Muñoz et al. (2022) described sociocultural barriers but lacked an exploration of emotional dimensions. This study complements and extends that literature by offering phenomenological insight into how women construct meaning through everyday interactions with mHealth. In contrast to more

outcome-driven studies (Yin et al., 2021; Krishnamurti et al. 2024), this study foregrounds the subjective voices of women, thereby contributing to a broader discourse on patient-centered digital health. The ambivalence experienced by users supports existing theories of digital health acceptance that emphasize emotional and contextual factors (Janols et al., 2022). Moreover, the emergence of trust through peer validation aligns with sociocultural theories of learning, which stress the importance of community influence in shaping behavior.

Beyond complementing existing work, these findings also challenge the assumption that digital health interventions are universally empowering. Instead, they demonstrate how poorly contextualized designs may reinforce feelings of isolation or burden. Theoretically, this underscores the need to integrate perspectives from critical digital health studies, which highlight how technology can reproduce social inequities if not carefully adapted to local realities (Lupton, 2019). By situating psychosocial experiences within this broader debate, the study emphasizes that effective digital maternal care requires a balance between technological innovation and culturally sensitive, human-centered design.

## **CONCLUSION**

This study explored the psychosocial experiences of pregnant women using mobile health (mHealth) applications in rural areas, addressing the gap in understanding how these technologies are lived and interpreted. The findings revealed four key themes—emotional ambivalence, digital isolation, cognitive fatigue, and evolving trust—that shape how women engage with mHealth during pregnancy. These insights offer a deeper understanding of the emotional and contextual factors that influence the adoption of digital maternal health tools, which have often been overlooked in prior outcome-driven research. By using a descriptive phenomenological approach, the study provides a human-centered perspective that emphasizes the importance of empathy, cultural sensitivity, and sustained support in mHealth implementation.

For policymakers, the findings suggest the need to design digital maternal health initiatives that are integrated into existing community health systems, ensuring that digital tools are accompanied by human facilitation from midwives or community health workers. Investment in rural digital infrastructure and literacy training should be prioritized to reduce digital isolation and cognitive burden among pregnant women. For developers, the study highlights the importance of incorporating emotional usability into application design—such as simplified language, culturally relevant content, interactive features that mimic empathetic communication, and peer-support functions that foster trust and social reassurance. Notifications and data input requirements should be streamlined to reduce fatigue, while adaptability to local contexts must guide interface design.

The results suggest that future mHealth strategies should prioritize emotional usability alongside technical functionality to ensure long-term engagement. Collaborations between policymakers, healthcare providers, and technology developers will be crucial in creating sustainable, inclusive, and culturally responsive digital maternal care ecosystems. Further research could expand these findings by exploring different populations, incorporating longitudinal designs, or integrating perspectives from healthcare professionals and family members.

## **CONFLICT OF INTEREST**

The authors declare that there is no conflict of interest.

## **REFERENCES**

Antelo, V. S., Szwarc, L., Paolino, M., Saimovici, D., Massaccesi, S., Viswanath, K., & Arrossi, S. (2022). A Counseling Mobile App to Reduce the Psychosocial Impact of Human Papillomavirus Testing: Formative Research Using a User-Centered Design Approach in a

- Low-Middle-Income Setting in Argentina. *JMIR Formative Research*, 6(1). Scopus. <https://doi.org/10.2196/32610>
- Atujuna, M., Simpson, N., Ngobeni, M., Monese, T., Giovenco, D., Pike, C., Figerova, Z., Visser, M., Biriotti, M., Kydd, A., & Bekker, L.-G. (2021). Khuluma: Using Participatory, Peer-Led and Digital Methods to Deliver Psychosocial Support to Young People Living With HIV in South Africa. *Frontiers in Reproductive Health*, 3. Scopus. <https://doi.org/10.3389/frph.2021.687677>
- Badawy, S. M., Abebe, K. Z., Reichman, C. A., Checo, G., Hamm, M. E., Stinson, J., Laloo, C., Carroll, P., Saraf, S. L., Gordeuk, V. R., Desai, P., Shah, N., Liles, D., Trimnell, C., & Jonassaint, C. R. (2021). Comparing the effectiveness of education versus digital cognitive behavioral therapy for adults with sickle cell disease: Protocol for the cognitive behavioral therapy and real-time pain management intervention for sickle cell via mobile applications (CaRISMA) study. *JMIR Research Protocols*, 10(5). Scopus. <https://doi.org/10.2196/29014>
- Ben-Zeev, D., Larsen, A., Attah, D. A., Obeng, K., Beaulieu, A., Asafo, S. M., Gavi, J. K., Kadakia, A., Sottie, E. Q., Ohene, S., Kola, L., Hallgren, K., Snyder, J., Collins, P. Y., & Ofori-Atta, A. (2024). Combining mHealth Technology and Pharmacotherapy to Improve Mental Health Outcomes and Reduce Human Rights Abuses in West Africa: Intervention Field Trial. *JMIR Mental Health*, 11(1). Scopus. <https://doi.org/10.2196/53096>
- Birrell, L., Furneaux-Bate, A., Debenham, J., Spallek, S., Newton, N., & Chapman, C. (2022). Development of a Peer Support Mobile App and Web-Based Lesson for Adolescent Mental Health (Mind Your Mate): User-Centered Design Approach. *JMIR Formative Research*, 6(5). Scopus. <https://doi.org/10.2196/36068>
- Brogly, C., Shoemaker, J. K., Lizotte, D. J., Kueper, J. K., & Bauer, M. (2021). A mobile app to identify lifestyle indicators related to undergraduate mental health (smart healthy campus): Observational app-based ecological momentary assessment. *JMIR Formative Research*, 5(10). Scopus. <https://doi.org/10.2196/29160>
- Browne, S., Kechadi, M.-T., O'Donnell, S., Dow, M., Tully, L., Doyle, G., & O'Malley, G. (2020). Mobile health apps in pediatric obesity treatment: Process outcomes from a feasibility study of a multicomponent intervention. *JMIR mHealth and uHealth*, 8(7). Scopus. <https://doi.org/10.2196/16925>
- Cyriac, J., Jenkins, S., Patten, C. A., Hayes, S. N., Jones, C., Cooper, L. A., & Brewer, L. C. (2021). Improvements in diet and physical activity-related psychosocial factors among African Americans using a mobile health lifestyle intervention to promote cardiovascular health: The FAITH! (fostering african american improvement in total health) app pilot study. *JMIR mHealth and uHealth*, 9(11). Scopus. <https://doi.org/10.2196/28024>
- Emert, S. E., Taylor, D. J., Gartenberg, D., Schade, M. M., Roberts, D. M., Nagy, S. M., Russell, M., Huskey, A., Mueller, M., Gamaldo, A., & Buxton, O. M. (2023). A non-pharmacological multi-modal therapy to improve sleep and cognition and reduce mild cognitive impairment risk: Design and methodology of a randomized clinical trial. *Contemporary Clinical Trials*, 132. Scopus. <https://doi.org/10.1016/j.cct.2023.107275>
- Fan, X., She, R., Liu, C., Zhong, H., Lau, J. T. F., Hao, C., Li, J., Hao, Y., Li, L., & Gu, J. (2020). Evaluation of smartphone APP-based case-management services among antiretroviral treatment-naïve HIV-positive men who have sex with men: A randomized controlled trial protocol. *BMC Public Health*, 20(1). Scopus. <https://doi.org/10.1186/s12889-020-8171-5>
- Giunti, G., Yrttiaho, T., Guardado-Medina, S., Sachinopoulou, A., Mylonopoulou, V., Fält, J., Paloniemi, B., Ryytty, M., Krüger, J., & Isomursu, M. (2025). Feasibility and usability evaluation of a gamified fatigue management mobile application for persons with multiple

- sclerosis in everyday life. *Multiple Sclerosis and Related Disorders*, 97. Scopus. <https://doi.org/10.1016/j.msard.2025.106379>
- Granhölm, E., Holden, J., Dwyer, K., Mikhael, T., Link, P., & Depp, C. (2020). Mobile-assisted cognitive behavioral therapy for negative symptoms: Open single-arm trial with schizophrenia patients. *JMIR Mental Health*, 7(12). Scopus. <https://doi.org/10.2196/24406>
- Hui, C. L.-M., Lam, B. S.-T., Wong, A. K.-H., Tao, T. J., Ho, E. C.-N., Suen, Y.-N., Lee, E. H.-M., Chan, S. K.-W., Chang, W.-C., Chong, C. S.-Y., Siu, C. M.-W., Lo, W. T.-L., & Chen, E. Y.-H. (2021). ReMind, a smartphone application for psychotic relapse prediction: A longitudinal study protocol. *Early Intervention in Psychiatry*, 15(6), 1659–1666. Scopus. <https://doi.org/10.1111/eip.13108>
- Iglesias, M., Sinha, C., Vempati, R., Grace, S. E., Roy, M., Chapman, W. C., & Rinaldi, M. L. (2023). Evaluating a Digital Mental Health Intervention (Wysa) for Workers' Compensation Claimants: Pilot Feasibility Study. *Journal of Occupational and Environmental Medicine*, 65(2), E93–E99. Scopus. <https://doi.org/10.1097/JOM.0000000000002762>
- Janols, R., Sandlund, M., Lindgren, H., & Pettersson, B. (2022). Older adults as designers of behavior change strategies to increase physical activity—Report of a participatory design process. *Frontiers in Public Health*, 10. Scopus. <https://doi.org/10.3389/fpubh.2022.988470>
- Johnson, B. (2014). *Educational research: Quantitative, qualitative, and mixed approaches* (Fifth edition). Sage Publications.
- King, V. L., Siegel, G., Priesmeyer, H. R., Siegel, L. H., & Potter, J. S. (2023). Development and Evaluation of a Digital App for Patient Self-Management of Opioid Use Disorder: Usability, Acceptability, and Utility Study. *JMIR Formative Research*, 8. Scopus. <https://doi.org/10.2196/48068>
- Krishnamurti, T., Rodriguez, S., Wilder, B., Gopalan, P., & Simhan, H. N. (2024). Predicting first time depression onset in pregnancy: Applying machine learning methods to patient-reported data. *Archives of Women's Mental Health*, 27(6), 1019–1031. Scopus. <https://doi.org/10.1007/s00737-024-01474-w>
- Lattie, E., Cohen, K. A., Winquist, N., & Mohr, D. C. (2020). Examining an app-based mental health self-care program, intellicare for college students: Single-arm pilot study. *JMIR Mental Health*, 7(10). Scopus. <https://doi.org/10.2196/21075>
- Mills, K. A. (2019). *Big data for qualitative research* (p. 78). Scopus. <https://doi.org/10.4324/9780429056413>
- Morcillo-Muñoz, Y., Sánchez-Guarnido, A. J., Calzón-Fernández, S., & Baena-Parejo, I. (2022). Multimodal Chronic Pain Therapy for Adults via Smartphone: Randomized Controlled Clinical Trial. *Journal of Medical Internet Research*, 24(5). Scopus. <https://doi.org/10.2196/36114>
- Morton, E., Nicholas, J., Yang, L., Lapadat, L., Barnes, S. J., Provencher, M. D., Depp, C., Chan, M., Kulur, R., & Michalak, E. E. (2022). Evaluating the quality, safety, and functionality of commonly used smartphone apps for bipolar disorder mood and sleep self-management. *International Journal of Bipolar Disorders*, 10(1). Scopus. <https://doi.org/10.1186/s40345-022-00256-6>
- Norden, M., Hofmann, A. G., Meier, M., Balzer, F., Wolf, O. T., Böttinger, E., & Drimalla, H. (2022). Inducing and Recording Acute Stress Responses on a Large Scale With the Digital Stress Test (DST): Development and Evaluation Study. *Journal of Medical Internet Research*, 24(7). Scopus. <https://doi.org/10.2196/32280>
- Pickard, A., Edwards, K., Farrow, C., Haycraft, E., & Blissett, J. (2025). Capturing Everyday Parental Feeding Practices and Eating Behaviors of 3- to 5-Year-Old Children With Avid Eating

- Behavior: Ecological Momentary Assessment Feasibility and Acceptability Study. *JMIR Formative Research*, 9. Scopus. <https://doi.org/10.2196/66807>
- Potzel, A. L., Gar, C., Seissler, J., & Lechner, A. (2021). A smartphone app (TRIANGLE) to change cardiometabolic risk behaviors in women following gestational diabetes mellitus: Intervention mapping approach. *JMIR mHealth and uHealth*, 9(5). Scopus. <https://doi.org/10.2196/26163>
- Psihogios, A. M., King-Dowling, S., Ohagan, B., Darabos, K., Maurer, L., Young, J., Fleisher, L., Barakat, L. P., Szalda, D., Hill-Kayser, C. E., & Schwartz, L. A. (2021). Contextual Predictors of Engagement in a Tailored mHealth Intervention for Adolescent and Young Adult Cancer Survivors. *Annals of Behavioral Medicine*, 55(12), 1220–1230. Scopus. <https://doi.org/10.1093/abm/kaab008>
- Rabbitts, J. A., Zhou, C., de la Vega, R., Aalfs, H., Murray, C. B., & Palermo, T. M. (2021). A digital health peri-operative cognitive-behavioral intervention to prevent transition from acute to chronic postsurgical pain in adolescents undergoing spinal fusion (SurgeryPalTM): Study protocol for a multisite randomized controlled trial. *Trials*, 22(1). Scopus. <https://doi.org/10.1186/s13063-021-05421-3>
- Rangraz Jeddi, F., Nabovati, E., Mobayen, M., Akbari, H., Feizkhah, A., Osuji, J., & Bagheri Toolaroud, P. (2023). Health care needs, eHealth literacy, use of mobile phone functionalities, and intention to use it for self-management purposes by informal caregivers of children with burns: A survey study. *BMC Medical Informatics and Decision Making*, 23(1). Scopus. <https://doi.org/10.1186/s12911-023-02334-w>
- Sauer, C., Zschäbitz, S., Krauss, J., Walle, T., Haag, G. M., Jäger, D., Hiller, K., Bugaj, T. J., Friederich, H.-C., & Maatouk, I. (2024). Electronic health intervention to manage symptoms of immunotherapy in patients with cancer (SOFIA): Results from a randomized controlled pilot trial. *Cancer*, 130(14), 2503–2514. Scopus. <https://doi.org/10.1002/cncr.35300>
- Sîrbu, V., David, O. A., Sanchez-Lopez, A., & Blanco, I. (2025). Comparative efficacy of PsyPills and OCAT mobile psychological interventions in reducing depressive, anxiety and stress symptoms: A blinded randomized clinical trial. *Journal of Affective Disorders*, 369, 945–953. Scopus. <https://doi.org/10.1016/j.jad.2024.10.052>
- Tatar, O., Abdel-Baki, A., Dyachenko, A., Bakouni, H., Bahreman, A., Tibbo, P. G., Crockford, D., Roy, M.-A., Copeland, J., Fischer, B., Lecomte, T., Côté, J., Ouellet-Plamondon, C., L'Heureux, S., Villeneuve, M., & Jutras-Aswad, D. (2023). Evaluating preferences for online psychological interventions to decrease cannabis use in young adults with psychosis: An observational study. *Psychiatry Research*, 326. Scopus. <https://doi.org/10.1016/j.psychres.2023.115276>
- Valle, C. G., Pinto, B. M., LaRose, J. G., Diamond, M., Horrell, L. N., Nezami, B. T., Hatley, K. E., Coffman, E. M., Polzien, K., Hales, D. P., Deal, A. M., Rini, C. M., Rosenstein, D. L., & Tate, D. F. (2021). Promoting physical activity in young adult cancer survivors using mHealth and adaptive tailored feedback strategies: Design of the Improving Physical Activity after Cancer Treatment (IMPACT) randomized controlled trial. *Contemporary Clinical Trials*, 103. Scopus. <https://doi.org/10.1016/j.cct.2021.106293>
- van der Storm, S. L., Bemelman, W. A., van Dieren, S., & Schijven, M. P. (2023). A personalized app to improve quality of life of patients with a stoma: A protocol for a multicentre randomized controlled trial. *Colorectal Disease*, 25(10), 2071–2077. Scopus. <https://doi.org/10.1111/codi.16694>
- van der Storm, S. L., Consten, E. C. J., Govaert, M. J. P. M., Tuynman, J. B., Oosterling, S. J., Grotenhuis, B. A., Smits, A. B., Marsman, H. A., van Rossem, C. C., van Duyn, E. B., de Nes, L. C. F., Verdaasdonk, E., de Vries Reilingh, T. S., Vening, W., Bemelman, W. A., & Schijven, M. P. (2024). Better stoma care using the Stoma App: Does it help? A first

- randomized double-blind clinical trial on the effect of mobile healthcare on quality of life in stoma patients. *Surgical Endoscopy*, 38(3), 1442–1453. Scopus. <https://doi.org/10.1007/s00464-023-10593-x>
- van der Storm, S. L., Hensen, N., & Schijven, M. P. (2023). Patient satisfaction with stoma care and their expectations on mobile apps for supportive care. *Colorectal Disease*, 25(9), 1852–1862. Scopus. <https://doi.org/10.1111/codi.16658>
- Wan, S. W., Chong, C. S., Toh, E.-L., Lim, S. H., Loi, C. T. T., Lew, Y. F. H., Chua, M. C. H., Jee, X. P., Liu, G., Zhu, L., Pikkarainen, M., & He, H.-G. (2021). A Theory-Based, Multidisciplinary Approach to Cocreate a Patient-Centric Digital Solution to Enhance Perioperative Health Outcomes among Colorectal Cancer Patients and Their Family Caregivers: Development and Evaluation Study. *Journal of Medical Internet Research*, 23(12). Scopus. <https://doi.org/10.2196/31917>
- Ward, R. E., Setiawan, I. M. A., Quinby, E., Fair, M., Ambadar, Z., Parmanto, B., & Dicianno, B. E. (2022). Mobile Health to Support Community-Integration of Individuals With Disabilities Using iMHere 2.0: Focus Group Study. *JMIR Human Factors*, 9(1). Scopus. <https://doi.org/10.2196/31376>
- Wong, R. S. M., Yu, E. Y. T., Wong, T. W.-L., Fung, C. S. C., Choi, C. S. Y., Or, C. K. L., Liu, K. S. N., Wong, C. K. H., Ip, P., & Lam, C. L. K. (2020). Development and pilot evaluation of a mobile app on parent-child exercises to improve physical activity and psychosocial outcomes of Hong Kong Chinese children. *BMC Public Health*, 20(1). Scopus. <https://doi.org/10.1186/s12889-020-09655-9>
- Yin, Y., Wang, H., Fan, C.-F., & Chen, H. (2021). Potential development of a mobile application for gout self-management: What support do patients need? *Patient Preference and Adherence*, 15, 2231–2238. Scopus. <https://doi.org/10.2147/PPA.S310689>
- Yoon, S., Goh, H., Nadarajan, G. D., Sung, S., Teo, I., Lee, J., Ong, M. E. H., Graves, N., & Teo, T. L. (2021). Perceptions of mobile health apps and features to support psychosocial well-being among frontline health care workers involved in the COVID-19 pandemic response: Qualitative study. *Journal of Medical Internet Research*, 23(5). Scopus. <https://doi.org/10.2196/26282>